

A Dissertation on
NON ALCOHOLIC FATTY LIVER DISEASE FIBROSIS SCORE
IN DIABETIC PATIENTS

Submitted to
THE TAMILNADU Dr. M.G.R MEDICAL UNIVERSITY
CHENNAI

*in partial fulfillment of the regulations
for the award of the degree of*

M.D BRANCH - I
GENERAL MEDICINE



GOVT.STANLEY MEDICAL COLLEGE & HOSPITAL
CHENNAI – TAMILNADU

MAY 2018

CERTIFICATE

This is to certify that this dissertation entitled “**NON ALCOHOLIC FATTY LIVER DISEASE FIBROSIS IN DIABETIC PATIENTS**” submitted by **DR.SINDHIYA JAYACHANDRAN** to the Tamilnadu Dr. M.G.R medical University is in partial fulfillment of the requirement of the award of **M.D DEGREE (BRANCH-I)** and is a bonafide research work carried out by her under direct supervision and guidance.

Signature of the Unit Chief

DR.P.VASANTHI.,M.D.,

Signature of HOD

DR.P.VASANTHI.,M.D.,

Signature of the dean

PROF.DR.S.PONNAMBALA NAMSHIVAYAM., M.D., D.A,D.N.B

DECLARATION

I solemnly declare that the dissertation entitled **“NON ALCOHOLIC FATTY LIVER DISEASE FIBROSIS IN DIABETIC PATIENTS”** was done by me at the Government Stanley Medical College and Hospital during **MARCH-SEPTEMBER 2017** under the guidance and supervision of **Prof.Dr.P.VASANTHI. M.D.** The dissertation is submitted to the Tamilnadu Dr.M.G.R Medical University towards the partial fulfillment of requirement for the award of M.D. Degree (Branch-I) in General Medicine.

Place:

Date :

DR.SINDHIYA JAYACHANDRAN

ACKNOWLEDGEMENT

I owe my thanks to **Prof. DR.S.PONNAMBALA NAMASIVAYAM,M.D, D.A,D.N.B** The Dean, Government Stanley Medical College, for allowing me to avail the facilities needed for my dissertation work.

I am grateful to **Prof. Dr.P.VASANTHI M.D.,** Professor and Head of the Department of General Medicine, Government Stanley Medical College for permitting me to do the study.

I am grateful to **Prof.DR.A.R.VENKATESHWARAN,M.D,D.M.,** Professor and Head of Department of General Medical Gastroenterology, Government Stanley Medical College for guiding me to do the study.

I express my gratitude to my unit Assistant Professors **Dr.NAMITHANARAYANAN M.D.,Dr.K.RAJA M.D.,** for their valuable suggestions and guidance.

I thank all my patients for letting me to examine them and cooperated to the procedures, patiently.

INSTITUTIONAL ETHICAL COMMITTEE,
STANLEY MEDICAL COLLEGE, CHENNAI-1

Title of the Work : Non Alcoholic Fatty liver disease fibrosis score in Diabetic Patients.

Principal Investigator : Dr. Sindhiya Jayachandran

Designation : PG MD (General Medicine)


Department : Department of General Medicine
Government Stanley Medical College,
Chennai-01

The request for an approval from the Institutional Ethical Committee (IEC) was considered on the IEC meeting held on 24.02.2017 at the Council Hall, Stanley Medical College, Chennai-1 at 2PM

The members of the Committee, the secretary and the Chairman are pleased to approve the proposed work mentioned above, submitted by the principal investigator.

The Principal investigator and their team are directed to adhere to the guidelines given below:

1. You should inform the IEC in case of changes in study procedure, site investigator investigation or guide or any other changes.
2. You should not deviate from the area of the work for which you applied for ethical clearance.
3. You should inform the IEC immediately, in case of any adverse events or serious adverse reaction.
4. You should abide to the rules and regulation of the institution(s).
5. You should complete the work within the specified period and if any extension of time is required, you should apply for permission again and do the work.
6. You should submit the summary of the work to the ethical committee on completion of the work.


MEMBER SECRETARY M15713
IEC-SMC, CHENNAI
MEMBER SECRETARY
ETHICAL COMMITTEE,
STANLEY MEDICAL COLLEGE
CHENNAI-600 001.

PLAGIARISM CERTIFICATE



Urkund Analysis Result

Analysed Document: DISSERTATION SINDHIYA (Autosaved)-1.docx (D31229099)
Submitted: 10/11/2017 8:22:00 PM
Submitted By: meyappansindhi@gmail.com
Significance: 1 %

Sources included in the report:

dissertation divakar1.docx (D31204219)

Instances where selected sources appear:

2

PLAGIARISM PERCENTAGE

URKUND

Document

Submitted

Submitted by

Receiver

Message

DISSERTATION SINDHIYA (Autosaved)-1.docx (D31229099)

2017-10-11 23:52 (+05:0-30)

Sindhiya jayachandran (meyappansindhi@gmail.com)

meyappansindhi.mgmu@analysis.urkund.com

Regarding plagiarism for dissertation submission [Show full message](#)

Sources

Highlights

Rank

Path/Filename

>

1

dissertation divakar1.docx

Alternative sources

Sources not used

1 Warnings

Reset

Export

Share

1% of this approx. 21 pages long document consists of text present in 1 sources.

A Dissertation on NON ALCOHOLIC FATTY LIVER DISEASE FIBROSIS SCORE IN DIABETIC PATIENTS
Submitted to THE TAMILNADU Dr. M.G.R MEDICAL UNIVERSITY CHENNAI

in partial fulfillment of the regulations for the award of the degree of

M.D BRANCH-II GENERAL MEDICINE

GOVT.STANLEY MEDICAL COLLEGE & HOSPITAL CHENNAI - TAMILNADU APRIL 2018

CERTIFICATE

This is to certify that this dissertation entitled "NON ALCOHOLIC FATTY LIVER DISEASE FIBROSIS IN DIABETIC PATIENTS" submitted by DR.SINDHIYA JAYACHANDRAN to the Tamilnadu Dr. M.G.R medical University is in partial fulfillment of the requirement of the award of M.D DEGREE (BRANCH-II) and is a bonafide research work carried out by her under direct supervision and guidance.

Signature of the Unit Chief Signature of HOD DR.P.VASANTHI.,M.D., DR.P.VASANTHI.,M.D.,

Signature of the dean PROF.DR.S.PONNAMBALA NAMSHIVAYAM.M.D.,

DECLARATION

CONTENTS

S NO.	PARTICULARS	PAGENO.
1	INTRODUCTION	1
2	AIMS OF THE STUDY	3
3	REVIEW OF LITERATURE	5
4	MATERIALS AND METHODS	38
5	OBSERVATION AND RESULTS	42
6	DISCUSSIONS	53
7	SUMMARY	56
8	TABLES AND FIGURES	58
9	CONCLUSIONS	74

ANNEXURE

I	MASTER CHART
II	BIBLIOGRAPHY
III	PROFORMA

INTRODUCTION

1. INTRODUCTION

Non alcoholic fatty liver disease is increasingly being recognised as a major cause of liver related morbidity and mortality, because of its potential to progress to cirrhosis and liver failure. NAFLD is deposition of fat in the liver of a non-alcoholic subject, a condition which may progress to end stage liver disease. (1)

NAFLD is an extremely common liver disease in the United States (USA), affecting approximately 20% of the adult population. In different countries, its prevalence is 10-24% of the general population. Amongst the obese persons, the prevalence rises to 57-74% and 25-75% among obese diabetics. Accordingly, NASH is considered as the third commonest cause of liver disease after alcohol abuse and hepatitis C. Patients with Diabetes mellitus have an increased risk of developing the spectrum of non alcoholic fatty liver disease (NAFLD) manifestations ranging from simple steatosis, non alcoholic steatohepatitis, hepatic fibrosis, cirrhosis and liver failure. Furthermore Diabetes mellitus with NAFLD have three times high mortality compared to non-diabetic NAFLD.

NAFLD fibrosis score is non invasive predictive score for hepatic fibrosis which includes six parameters and categorised into three groups based on the scores. F0-F2 = <-1.455 , Intermediate = $-1.455-0.675$, F3-F4 = >0.675 .

AIM OF THE STUDY

2.AIM OF THE STUDY

1. To categorize the Type 2 diabetic patients into four groups-obese, non-obese, ultrasonographic presence and absence of hepatic steatosis.
2. To analyse the risk of hepatic fibrosis in these patients using NAFLD FIBROSIS SCORE.
- 3.To correlate the stage of fibrosis by fibro scan in group of patients with high NAFLD score.

REVIEW OF LITERATURE

3.REVIEW OF LITERATURE

Nonalcoholic fatty liver disease (NAFLD) is a broader term with a spectrum including patients with simple steatosis, steatohepatitis that can progress to cirrhosis liver and even hepatocellular carcinoma.(1)(2). In 1980, Ludwig et al(3) described the term NASH as a form of liver injury that was histologically consistent with alcoholic hepatitis but occurred in obese, diabetic females, who denied alcohol use. The prevalence of NAFLD and NASH in the general population is estimated to be between 10%-24% and between 1%- 5%, respectively. There is a direct correlation between body mass index (BMI) and prevalence and severity of NAFLD. The prevalence of NAFLD increases to 57.5% to 74% in obese persons and 90% in morbidly obese persons.(4)(5)(6).In India prevalence of NAFLD is around 9% to 32% of general population.(7)(8)

NAFLD not only affects the liver, it is a multisystem disease affecting extra-hepatic organs and regulatory pathways. It increases risk of type 2 diabetes mellitus, cardiovascular and cardiac diseases, chronic kidney disease and causal link in sleep apnea, colorectal cancers, osteoporosis, psoriasis and various endocrinopathies like polycystic ovary syndrome.

Most patients with NAFLD are asymptomatic and may have elevation of liver enzymes which is noted incidentally in biochemical investigations. Alanine amino transferase and aspartate amino transferase are usually elevated, but these levels do not reliably correlate with hepatic injury, inflammation or cirrhosis in NAFLD.

3.1.NATURAL HISTORY OF NAFLD:

About 90% of Patients with **NAFLD** have simple steatosis and therefore no increase in mortality (9)(10), approximately 10-30% patients have NASH (Non Alcoholic Steatohepatitis) which is associated with hepatocellular injury and inflammation. 25-40% of patients with NASH will develop progressive fibrosis and cirrhosis and end stage liver disease.(11)(12)(13).Recent data states that 28% of patients have progression on histological examination, 59% have no change, and 13% may have improvement. Matteoni et al retrospectively determined the histological and/or clinical outcome of 98 patients with the whole spectrum of NAFLD from simple steatosis through NASH to cirrhosis.

After a median 8-year follow up, 25% of individuals with evidence of hepatocyte necrosis with or without Mallory's hyaline or fibrosis, either already had cirrhosis on index biopsy or progressed to cirrhosis. This compared with only 3.4% of patients with simple steatosis with or

without non-specific inflammatory changes. These observations indicate that NAFLD patients without NASH have a benign prognosis.(14)

The development of cirrhosis due to NASH is associated with a poor prognosis. 10-year mortality rate is 20% for subjects with Child-Pugh A disease and 45% will decompensate within 10 years of diagnosis.(15)Patients with cirrhosis due to NASH are at significant risk of developing hepatocellular carcinoma.(16)

3.2.RISK FACTORS OF NAFLD:

3.2.1.AGE:

Higher risk is associated with increasing age of the patient.

3.2.2.METABOLIC SYNDROME:

Metabolic syndrome is independent predictor of fibrosis.70-90% of patients with NAFLD have metabolic syndrome. Insulin resistance is a key mediator between NAFLD and metabolic syndrome (17).Any three of the following features is diagnosed as metabolic syndrome.

- a. Central obesity – Waist circumference >94cm for men and >80cm for women.
- b. Impaired fasting glucose - >5.6 mmol/L or on treatment.
- c. Hypertriglyceridemia - >1.7 mmol/l or on treatment

- d. Low HDL cholesterol - <1.0 mmol/l for men or on treatment
<1.3 mmol/l for women or on treatment
- e. Hypertension - >135/85 mm Hg or on treatment

3.2.3.GENDER:

NAFLD is more common in men compared to women, but progression to advanced fibrosis is higher in women compared to men.(18)(19)

3.2.4.ETHNICITY:

NAFLD risk is higher in Hispanics population and lower in black population.(20)

3.2.5.DIETARY FACTORS:

Diet containing high cholesterol and saturated fats, high fructose intake, low carbohydrates increases the risk of NAFLD(21)(22).

Caffeine may be protective in some patients.(23)

3.2.6.OBSTRUCTIVE SLEEP APNOEA:

It increases the risk of NAFLD.(24)(25)

3.2.7. GENETIC FACTORS:

Patatin –like phospholipase domain- containing 3 (PNPLA3).
(25)(26)(27) PNPLA3 lysophosphatidic acid acyltransferase activity could also contribute to altered plasma triacylglycerol composition and concentration.

3.2.8.NUTRITIONAL FACTORS:

Obesity, rapid weight loss, total parenteral nutrition, prolonged starvation are attributed as causal factors in NAFLD.

3.2.9.DRUGS:

Corticosteroids, Amiodarone, Bleomycine, Tetracycline, Methotrexate, Perhexiline, Diltiazem, nifedipine , Tamoxiphen.

3.2.10.CHEMICALS:

Hydrocarbons and yellow phosphorous are associated with fatty liver.

3.2.112.SURGERY:

Surgeries like Jejunoileal bypass,extensive small bowel loss, gastropexy are associated with fatty liver.

3.3.ETIOPATHOGENESIS OF NAFLD/NASH:

The pathogenesis of the NASH is not completely understood. The conditions associated with NAFLD include nutritional abnormalities, metabolic disorders, drugs, chemicals and surgery as mentioned above. Initially there is steatosis which later, after a series of events, may progress on to inflammation and fibrosis and finally to cirrhosis and end stage liver disease.(28)(29)(30) NASH may be considered as a two hit process, first hit is accumulation of the fat and the second hit is hepatocellular injury in the fatty liver .(30)The first hit of steatosis occurs because of the imbalance between the fatty acid uptake, its oxidation, esterification and export as very low-density lipoprotein (VLDL) from the hepatocytes. Insulin resistance is the most important key factor(28)((30)(31)which leads to increase in lipolysis and increased uptake of fatty acids by hepatocytes.

Hyperinsulinemia which occurs as result of insulin resistance increases the intrahepatocytic fatty acids by increasing the glycolysis and decreasing the apolipoprotein B-100 and resulting in decreased export as VLDL. The end result is the increase in fatty acids and triglycerides in the hepatocytes leading to steatosis or fatty liver.

The free fatty acids in the hepatocytes acts as reactive oxygen species and increases the mitochondrial β -oxidation and cytochrome P-450 4A and cytochrome P450 2E1 levels. The mitochondrial oxidative stress leads to the second hit from steatosis to steatohepatitis and fibrosis by three main mechanisms, namely (i) lipid peroxidation, (ii) cytokine induction, and (iii) Faslig and induction.(28)(30)

(i). Lipid peroxidation causes oxidative destruction of polyunsaturated fatty acids of cellular membranes. The cytotoxic products which are released due to lipid peroxidation impair cellular functions including nucleotide and protein synthesis leading to cell death, formation of Mallory hyaline, promoting tissue inflammation, activation of stellate cells and collagen synthesis.(28)

(ii). Cytokines like IL-1, IL-6 IL-8 and tumor necrosis factor- α (TNF- α) plays an important role in the pathogenesis of liver injury in patients with NASH.(31)

IL-1, IL-6, and IL-8 are pro-inflammatory cytokines and IL-10 and IL-12 are anti-inflammatory cytokines. TNF- α , TGF- β , interleukin-8 and cause chemotaxis, formation of Mallory hyaline and synthesis of collagen by activated stellate cells.

TNF- α downregulates insulin-induced phosphorylation of insulin receptor substrate 1 and reduces the expression of the insulin-dependent glucose transport molecule Glut 4 and thus contributes towards insulin resistance which is thought to be the major mechanism in the pathogenesis of NASH.(31)(32)

TNF- α is derived from adipose tissue in the absence of active infections or inflammatory conditions. Plasma levels of TNF- α also correlates with body fat mass and is associated with insulin resistance. (37) TNF- α knockout mouse also fails to develop insulin resistance after induction of obesity suggests its crucial role in the pathogenesis of insulin resistance.(38) The oxidative stress leads to the I κ B activation and release of TNF- α and insulin resistance, which ultimately leads to steatohepatitis.

(iii). Finally the expression of Fas ligand due to oxidative stress leads to fractional killing by interaction with Fas on other hepatocytes.

Other factors that are involved in the pathogenesis of NASH include serum and liver iron,(33)(34) leptin,(35) bacterial overgrowth, (36)(37)(38) Saturation of mitochondrial β -oxidation leads to peroxisomal oxidation and generation of hydrogen peroxide, which in the

presence of increased iron is converted to hydroxyl radicals, thus adding to the oxidative stress and further injury.(39)

Leptin, a product of obesity gene regulates the food intake and body composition through a central feed back mechanism and is proposed to be a key pathophysiological factor for obesity. Leptin leads to hepatic steatosis by promoting insulin resistance or by modulating insulin signaling in hepatocytes.(34)

Bacterial overgrowth in the gut can cause liver injury by causing endotoxemia and release of cytokines. The mechanisms by which intestinal bacteria may increase hepatic oxidative stress include increased endogenous production of ethanol and by direct activation of inflammatory cytokines in luminal epithelial cells, and liver macrophages or both .(35)




FIG 1: The factors like visceral ectopic fat accumulation, adipose tissue inflammation, type 2 diabetes and intestinal dysbiosis which promote development of progressive liver disease in NAFLD.(Non Alcoholic Fatty Liver Disease)

3.4.NAFLD AND DIABETES MELLITUS:

When imbalance occurs between energy intake and energy expenditure, or an intrinsic problem with storing excess energy as lipid (triacylglycerol) in adipose tissue depots, lipid occurs in other organs throughout the body. Ectopic fat accumulation is defined as lipid accumulation in tissues, organs not designed to accumulate e.g., liver or omentum other than adipose tissue.(40)

NAFLD is an ectopic fat accumulation which is associated with increased secretion of hepatokines (41), increased gluconeogenesis, decreased glycogen synthesis and inhibition of insulin signaling (42)(43). Adipose tissue inflammation is crucial in NAFLD pathogenesis and evidence suggesting that dysbiosis of the gut microbiota plays a major key role in development and progression of NAFLD. This is mainly because of the increased intestinal absorption of bacterial products, such as short-chain fatty acids (e.g., butyrate, propionate and acetate), lipopolysaccharide and endotoxins.

Mechanisms involved in the pathogenesis of insulin resistance and inflammation in NAFLD is explained briefly. Although obesity is strongly associated with hepatic steatosis, excess body fat accumulation is not 'sine qua non' for developing NAFLD. In fact, patients with lipodystrophy have marked insulin resistance and commonly develop hepatic steatosis

and T2DM, strongly suggesting that it is not body fat mass per se that is important, but it is adipose tissue dysfunction that is a key contributor to the pathogenesis of NAFLD (46).

Specifically, increased free fatty acid (FFA) fluxes from the adipose tissue pool increase the availability of long-chain fatty acyl-CoAs for hepatic lipid accumulation, particularly in physically inactive individuals(44), and evidence is accumulating that hepatic lipid accumulation is capable of causing hepatic/peripheral insulin resistance and promoting hepatic inflammation (40)(44).Expansion of peripheral adipose depots provides buffering capacity that may protect the liver from the excessive FFA fluxes that promote hepatic lipid accumulation. Within hepatocytes, long-chain fatty acids are esterified with glycerol-3-phosphateto form mono-acylglycerols, di-acylglycerols and tri-acylglycerols.

Lipid synthesis is very important in causing ‘resistance’within the hepatic insulin signaling pathway (47), promoting hepatic inflammation (48) and increasing risk ofprogressive liver disease that occurs with NASH. In the liver ceramides can accumulate into the cells via three main routes: 1) the hydrolysis of the membrane phospholipid sphingomyelin, which is coordinated by the enzyme sphingomyelinase; 2) de novo synthesis from long chain fatty acids such as palmitate and

serine; and 3) a 'salvage' pathway that utilizes sphingosine and forms ceramide (54)(55). Ceramide plays an important role in causing insulin resistance.(55)

Hepatic lipids that are not esterified also induce endoplasmic reticulum stress, leading to the activation of c-Jun N-terminal kinases and NF- κ B (55), which are two major regulators of inflammatory pathways that also inhibit phosphorylation of insulin receptor substrate-1 (IRS-1) (56), potentially aggravating hepatic insulin resistance and increasing intra-hepatic cytokine production. Synthesis of lipids such as diacylglycerol (DAGs) is intimately related to inflammatory pathways, and DAGs may also contribute to hepatic production of inflammatory cytokines [e.g., tumour necrosis factor- α (TNF- α), interleukin-6 (IL-6)], and procoagulant factors [e.g., factor VIII, plasminogen activator inhibitor-1 (PAI-1)].

Intestinal microbiota dysbiosis may affect other hepatic lipid pathways, such as those involving bile acid metabolism, consequently increasing hepatic inflammation and fibrosis, and resulting in an increased risk of developing cirrhosis and HCC.

Till date, it is unclear whether improvements in NAFLD may ameliorate risk of T2DM or improve glycaemic control in people with

NAFLD who have developed T2DM, but it is plausible that resolution of liver fat and improvements in liver lipid metabolism might modify the risk of T2DM via a liver-specific effect. Such a liver-specific effect could be mediated by alteration in the secretion of multiple hepatokines or inflammatory cytokines that influence risk of diabetes. In NAFLD, secretion of diabetogenic hepatokines, such as retinol binding protein (RB)-4, fetuin-A, fibroblast growth factor (FGF)-21; or inflammatory biomarkers such as C-reactive protein (CRP), TNF- α and IL-6 (57) may directly affect risk of incident T2DM by adversely affecting hepatic gluconeogenesis, glycogen synthesis and insulin signaling (58).




FIG 2 : Mechanism of insulin resistance, adipose tissue inflammation and role of hepatokines, adipokines in NAFLD.

3.5.NAFLD AND METABOLIC SYNDROME:

Metabolic syndrome is characterized by the presence of insulin resistance in association with other metabolic abnormalities like obesity, diabetes, hypertriglyceridemia and hypertension. Common associations of NAFLD, such as hypertension, hyperuricemia, and polycystic ovary syndrome are also common to metabolic syndrome.

The chances of individual having NAFLD and NASH increase with increasing body weight, with 70-80% of obese individuals having NAFLD and 15-20% having NASH. Conversely 30-100% of NASH patients have obesity.

Truncal obesity is more commonly associated with NASH, also with causal link to diabetes and hypertension. In morbidly obese patients, the risk of liver disease progressively increases with the number of features of metabolic syndrome.(59)(60)Metabolic syndrome was present in 22% of Indian patients with NASH.(61)

3.6.NAFLD AND CARDIOVASCULAR DISEASES:

Non alcoholic fatty liver disease is associated with increased risk and morbidity for cardiovascular diseases and risk further increases in Type 2 diabetes mellitus. In addition patients with non alcoholic steatohepatitis have further increased risk of cardiovascular diseases.

Myocardial metabolism is primarily affected in NAFLD. By cardiac magnetic resonance imaging (MRI), Perseghin et al. reported that nonobese, nondiabetic, normotensive, young individuals with NAFLD had impaired myocardial energy metabolism (i.e., a lower phosphocreatine/adenosine triphosphate ratio, as measured by ³¹P-magnetic resonance spectroscopy and excessive fat accumulation in the epicardial area compared with control subjects without NAFLD. Interestingly, these myocardial metabolic alterations were detected despite normal left ventricular (LV) morphological features and systolic and diastolic functions (62). Lautamaki et al.(63)and Rijzewijk et al. (64)found that T2DM patients with higher intra-hepatic fat content on ¹H-MRS had increased myocardial insulin resistance and decreased myocardial perfusion compared with those with lower intra-hepatic fat content, additionally, myocardial insulin resistance was more severe among those with higher intra-hepatic fat content, those with higher intra-hepatic fat content had significantly higher myocardial fat content.

Interestingly, in this study cardiac steatosis was a strong predictor of LV diastolic dysfunction.(65) NAFLD with increased liver transaminases is an independent risk factor for atrial fibrillation in the Framingham Heart Study cohort (66).

Asimilar link was found between elevated serum liver enzymes and AF risk was shown in a larger prospective community-based study of 9333 subjects with a follow up for 12 years (67). NAFLD is also independently linked with prolonged QTc interval which is a powerful predictor of ventricular arrhythmias and sudden cardiac death and the presence of progressive aortic valve sclerosis. All these factors contributes to increased cardiovascular risk in NAFLD patients.(68)(69)

3.7.NAFLD AND CHRONIC KIDNEY DISEASE:-

The prevalence of chronic kidney disease is increased among patients with non alcoholic fatty liver disease with or without diabetes mellitus. The estimated glomerulation filtration rate and overt proteinuria is measured in patients with chronic kidney disease.(70)The severity of NAFLD varied upon the stages of chronic kidney disease. The patients had fatty kidney and increased renal sinus fat volume caused structural and functional derangements in the kidneys.(71)




FIGURE 3: The link between NAFLD and cardiovascular and kidney structural and functional changes.

3.8. CLINICAL FEATURES:

1.SYMPTOMS:-

- Asymptomatic (30-40%)
- Right upper subcostal pain or discomfort (30-40%)
- Fatigue (<10%)
- Pedal edema
- Gastrointestinal bleeding
- Abdominal distension

2. SIGNS:-

- Normal examination
- Increased body mass index (BMI)
- Increased waist circumference (truncal obesity)
- Stigmata of chronic disease if cirrhosis is present.
- Lipomatosis/lipoatrophy /lipodystrophy
- Hepatomegaly (50-70%)
- Splenomegaly
- Ascites

3.9. INVESTIGATIONS:

3.9.1. BIOCHEMICAL :

- i. Raised aspartate aminotransferase (AST) and alanine aminotransferase (ALT) - up to 4-5 times elevation
- ii. Alkaline phosphatase - up to 2 times elevation
- iii. AST/ALT - majority <1 ; >1 may indicate cirrhosis
- iv. Bilirubin - elevated in late stage
- v. Albumin - decreased in late stage
- vi. Prothrombin time - prolonged in late stage
- vii. Serum markers of iron overload - ~25% but do not indicate hemochromatosis.
- viii. Anti-nuclear antibody - one-third

3.9.2. FIBROSIS SCORES:

1.NAFLD FIBROSIS SCORE:

Fibrosis staging is important in all patients with NAFLD to identify patients with advanced hepatic fibrosis at risk of liver-related complications. Hepatocellular dysfunction and portal hypertension result from advancing hepatic fibrosis. This can be noticed in blood investigations such as liver function tests (low albumin), full blood count (thrombocytopenia) and coagulation profile (prolonged prothrombin time). These tests give an indirect measure of fibrosis and act as non-

invasive markers of fibrosis as they are less cost and are performed in all patients with hepatic disease. With increasing hepatic fibrosis, the ALT typically falls and the AST remains stable or rises and as a result, the AAR (aspartate and alanine amino transferase ratio) increases and can be an useful simple method of identifying patients with advanced fibrosis. Previous studies identified a cut-off >1 for the AAR as a diagnostic test for cirrhosis. (72)(73)

2.BARD SCORE:

The BARD score is a simple test using the body mass index ≥ 28 – 1 point, AST/ALT ratio ≥ 0.8 – 2 points and presence of type 2 diabetes mellitus – 1 point. The score ranges from 0 to 4 points. A score <2 has excellent negative predictive value of 95–97%, which excludes advanced hepatic fibrosis.

However, in a typical NAFLD cohort, a large proportion of patients with mild disease have a score of ≥ 2 because of diabetes limiting its utility in clinical practice. (74)

3.FIB-4 SCORE:

The FIB-4 score although derived in patients with HIV and HEPATITIS B coinfection, appears to be one of the most essential noninvasive tests for diagnosing advanced fibrosis in NAFLD. For stage

3–4 fibrosis, a score <1.3 has a 90% NPV and a score >2.67 has an 80% PPV, with 72% of patients scoring below 1.3 or above 2..Other studies have proved that the FIB-4 score is considered slightly better than other non-invasive tests in diagnosing advanced fibrosis in NAFLD, including in subjects with normal range ALT levels.(75)

4.CK-18:

Levels of cytokeratin-18 (CK-18) fragments correlate with the magnitude of Hepatocyte apoptosis and predict the presence of NASH(non alcoholic steatohepatitis).Cytokeratins are proteins of keratin containing intermediate filaments found in intracytoplasmic cytoskeleton of epithelial tissue. Mallory body composed of abnormally phosphorylated cross linked keratins like cytokeratin 8 and 18.

5.HAIR SCORE:-

1. Alanine transaminase (ALT) > 40 IU/l
2. Insulin resistance (IR) index > 5
3. Hypertension

Presence of 2 or all 3 factors predict NASH.

6. BAAT SCORE:-

1. Age > 50 yrs.
2. Body mass index (BMI) > 28 kg/m².

3. Triglyceride > 1.7 mmol/l.

4. Alanine aminotransferase (ALT) > 2-fold rise.

Presence of none or only 1 factor rules out the possibility of fibrosis or cirrhosis.

7.OTHER NON INVASIVE TESTS:

The Enhanced Liver Fibrosis (ELF) test is a commercial panel of markers of matrix turnover: tissue inhibitor of matrix metalloproteinase 1 (TIMP1), hyaluronic acid and PIIINP.(76) Fibrotest (FT) is a commercial tool of biochemical markers of fibrosis that is prevalently used in France. In NAFLD, FT can diagnose advanced hepatic fibrosis with modest accuracy .Using a FT cut-off of 0.30 gives a 90% NPV for advanced fibrosis (sensitivity 77%), and a FT cut-off of 0.70 had a 73% PPV for advanced fibrosis (specificity 98%).Other scores include Palekar's score,Gholam score,Nippon score and BAAT score.(77)

3.9.3.IMAGING:

- **Ultrasound :**


Findings of hepatic steatosis in ultrasound are increased echogenicity and coarsened echotexture of the .If steatohepatitis has progressed to cirrhosis, a nodular liver surface may be present in addition to other fibrotic changes.

- **Fibro scan:-**


Fibrotic livers have less elasticity due to the fibrous tissue deposition in the hepatic parenchyma. TE (Fibroscan) gives a 'liver stiffness measurement'(LSM) using pulsed-echo ultrasound as a surrogate marker of fibrosis. The LSM correlates well with the degree of liver fibrosis in a wide range of liver diseases, including NAFLD.(78)




However, there are multiple drawbacks in using TE in NAFLD. Invalid results appear in older patients (>52 years) and those with central obesity (BMI >35 kg/m²) or type 2 diabetes. The Fibroscan XL probe has been developed for obese patients but is associated with fewer LSM failures.





Liver stiffness cut-offs in chronic liver diseases



Matavir	F0-F1	F2	F3	F4
Fibrosis	Mild	Sign	Severe	Cirrhosis
				

- **Computed tomography/Magnetic resonance imaging**

A. CT scan findings in NAFLD:

- Picking up of focal areas of fatty infiltration.
- Mean CT Hounsfield unit in liver(diffuse hypoattenuation) less than that inspleen helps in diagnosis.

B. MRI findings in NAFLD:

- Phase contrast imaging gives adequate information about fatty infiltration of liver giving very good quantitativeassessment of status of disease.
- Useful for excluding fatty infiltration.
- On T1-weighted images, there is loss of intensity in focal areas of deposition of fat. So,in early stage of disease ,small lesions are readily identifiedon MRI
- It is more sensitive and specific compared to USG or CTscan, but is expensive.

C. Radionuclide scanning (scintigraphy) studies:


- With technetium-99m sulphur colloid scanning, focal areas of deposition of fat appear as fillingdefects.
- Radio-xenon has a very high affinity for fat and itremains bound to fat and retained in fat. This gives qualitative as well as quantitative assessment offat deposition in liver.

- **ARFI:**

The non-invasive assessment of hepatic fibrosis is acoustic radiation force impulse (ARFI). This technique uses conventional B-mode ultrasonography to generate an ultrasonic pulse and measurement of the response of the liver tissue as shear wave velocity. The degree of hepatic fibrosis correlates with median velocity measured by ARFI.(79)

3.9.4.LIVER BIOPSY:

Performing a liver biopsy on every patient with suspected NAFLD remains a controversial area in practice, and clearly is not a practical consideration as a “screening” tool. The general indications for performing a liver biopsy in patients with NAFLD are to confirm or exclude the diagnosis, diagnose other liver diseases, and to determine amounts of damage to the liver for treatment and prognosis. The last includes necroinflammatory activity, which is potentially reversible, and collagen deposition with varying degrees of remodeling, which is potentially less reversible. The major limitation of liver biopsy is the invasive nature of the procedure.(80) Though considered “minimal”, liver biopsy is an invasive procedure and can have complications even in the ideal clinical conditions, including pain, minor and major bleeding (0.3%).(81)



A: Mixed large and small droplet steatosis, single droplet, with nucleus pushed to one side,

B: Microvesicular steatosis, nuclei in the center with foamy cytoplasm, and mega mitochondria

C: Ballooned hepatocytes with flocculent cytoplasm.

D: Loss of cytoplasmic expression of keratin 8/18 in ballooned hepatocytes,

E: Mallory-Denk body

F: Mallory-Denk body in blue-green color and dense perisinusoidal fibrosis.

G: Portal lipogranuloma.

H: Mallory-Denk bodies and satellitosis.

I: Delicate perisinusoidal fibrosis.

J: Bridging fibrosis.

TREATMENT OF NON ALCOHOLIC FATTY LIVER DISEASE:

Treatment is aimed at correcting the risk factors for NAFLD, such as, medical control of hyperglycemia, and use of lipid lowering agents for hypertriglyceridemia, and lifestyle modification and pharmacological agents to improve insulin sensitivity.

INSULIN SENSITIZERS:-

Thiazolidinediones are a class of drugs that stimulate peroxisome proliferator activated receptors (PPAR). An insulin-sensitizing medication, troglitazone (a thiazolidinedione) is used.

LIPID LOWERING AGENTS:

Hypertriglyceridemia is associated with NAFLD. Clofibrate and gemfibrozil can be used.

PHARMACOLOGICAL THERAPY OFFERING HEPATOCYTE PROTECTION:

Ursodeoxycholic acid (UDCA), and the anti-oxidants, betaine and vitamin E, have peer-reviewed published data. Other drugs, e.g., lecithin, β -carotene, selenium, and Nacetylcysteine.

UDCA:-

This hydrophilic bile acid with hepatoprotective properties was first evaluated in a pilot study, where it was associated with both improved liver enzyme levels and a decrease in hepatic steatosis.

Vitamin E (α -Tocopherol):-

The observation that vitamin E decreases oxidative stress provides a rationale For its use in patients with NASH. One uncontrolled trial in children with NASH showed that supplementation with vitamin E (400 to 1200 IU daily) was associated with a significant decline in serum aminotransferases.

There was a significant improvement in fibrosis scores in the NASH patients receiving vitamins compared to baseline but no improvement in necroinflammation or ALT.

Betaine:-

Betaine, a normal component of the metabolic cycle of methionine, is a precursor of *S*-adenosylmethionine, a hepatoprotective factor.

Liver Transplantation is the choice of treatment in end stage liver disease.

MATERIALS AND METHODS

4.MATERIALS AND METHODS

This study was carried out in Govt Stanley Medical college Hospital, Chennai. This is a facility based observational study involved the patients who attended medicine and diabetology outpatient departments and as well as inpatients during the study period of March 2017-September 2017.

STUDY DESIGN:OBSERVATIONAL STUDY

DATE OF APPROVAL BY ETHICAL COMMITTEE:FEBRUARY 2017

PLACE OF STUDY:

DEPARTMENT OF INTERNAL MEDICINE,GOVERNMENT STANLEY MEDICAL GENERAL HOSPITAL,CHENNAI.

SELECTION OF CASES:

Diabetic outpatients and inpatients attending our institute.

SAMPLE SIZE:

A total of 100 patients of diabetes mellitus who attended the outpatients department and admitted in medicine wards.

4.1 .INCLUSION CRITERIA:

1.All patients diagnosed as Type 2 Diabetes Mellitus by WHO criteria.

2.Age more than 18 years.

4.2.EXCLUSION CRITERIA:

1.Patients less than 18 years and more than 85 years.

2.Patients with history of alcohol intake more than 30 grams/day in males and more than 20 grams/day in females.

3.Patients with history of jaundice,Hepatitis B and C infection.

4.Patients with history of following drug intake such as steroids, synthetic Oestrogens,calcium channel blockers, amiodarone, valproic acid, heparin, antiviral agents.

4.3.METHODOLOGY:

Patients were made to understand in their local language and informed consent was obtained before collecting data and subjecting to investigations.

Patients were divided into four groups -obese group and non obese group according to WHO BMI criteria.BMI calculated by Quetelet index – Weight in kilograms / height in m²Ultrasonographic presence and absence of hepatic fibrosis.

- OBESE GROUP- BMI > 30kg/m²
- NON OBESE GROUP – BMI <29.9 kg/m².

ASSESMENT OF PARAMETERS:

- 1.Age
- 2.Diabetes mellitus
- 3.Body mass index (BMI)
- 4.Aspartate aminotransferase level.
- 5.Alanine aminotransferase level.
- 6.Platelet count.

FORMULA OF NAFLD FIBROSIS SCORE:

$$-1.675 + 0.037 * \text{AGE (years)} + 0.094 * \text{BMI (kg/m}^2\text{)} + 1.13 \\ * \text{IFG/ DIABETES (yes =1,no=0)} + 0.99 * \text{AST/ALT ratio} - \\ 0.013 * \text{platelet (* 10}^9\text{/l)} - 0.66 * \text{albumin (g/dl)}.$$

EXPLANATION OF RESULT:

- NAFLD SCORE <-1-455 = F0 –F2 (LOW RISK)
- NAFLD SCORE -1.455- 0.675 = indeterminate score
- NAFLD SCORE >0.675 = F3- F4. (HIGH RISK)

OBSERVATION & CALCULATION

5.OBSERVATION

100patients of Type 2 diabetes mellitus who attended general medicine and diabetology outpatient and inpatient departments in government stanley medical college during the period of march 2017 to september 2017 was studied and data collected and interpreted.

5.1.PATIENT CHARACTERISTICS:

5.1.1.PATIENT PROFILE:

Out of 100 patients,40 patients were male and 60 patients were females and patients were grouped into obese, non-obese, ultrasonographic presence of hepatic steatosis and ultrasonographic absence of hepatic steatosis.

Out of 100 patients, 57 patients (57%) were in obese group and 43 patients (43%)in non- obese group.

34 patients (59.6%) had ultrasonographic evidence of hepatic steatosis in the obese group.3 patients (6.9%) had ultrasonographic evidence of hepatic steatosis in non obese group.

23 patients (40.4%) had no ultrasonographic evidence of hepatic steatosis in obese group. 40 patients (93%) had no ultrasonographic evidence of hepatic steatosis in non-obese group.

5.2.PARAMETERS STUDIED:

5.2.1.NAFLD FIBROSIS SCORE:

Non alcoholic fatty liver disease fibrosis score was applied to all patients in both obese and non obese group and categorised into high risk, intermediate risk and low risk.

Accordingly 20 patients(35%) had high risk in obese group, 1 patient (0.2%) had high risk in non obese group.

21 patients (36%) had intermediate risk in obese group and 6 patients (13.9%) had intermediate risk in non obese group.

16 patients (28%) had low risk in obese group and 36 patients (83.7%) had low risk in non obese group.

5.2.2.FIBRO SCAN:

Non invasive fibroscan/ elastography was done to calculate the liver stiffness in patients with non alcoholic fatty liver disease and high NAFLD score.

Accordingly 20 patients of obese group had high risk of hepatic steatosis and fibro scan was done for the patients.

4 patients (20 %) had F0-F1 stage of fibrosis in fibro scan(according to METAVIR staging)

13 patients (65%) had F2-F3 stage of fibrosis.

3 patients (15%) had F4 stage of fibrosis.

TABLE 1: AGE RANGE OF STUDY POPULATION

Agerange


	Frequency	Percent
Upto 35 yrs	6	6.0
36 - 45 yrs	27	27.0
46 - 55 yrs	32	32.0
56 - 65 yrs	26	26.0
Above 65 yrs	9	9.0
Total	100	100.0

TABLE 2: STATISTICS OF AGE

AGE

N	Valid	100
	Mean	50.83
	Median	50.00
	Std. Deviation	10.431
	Range	48
	Minimum	26
	Maximum	74

FIGURE 1: PIE CHART REPRESENTING THE PERCENTAGE OF OBESE AND NON OBESE GROUP IN STUDY POPULATION



**FIGURE 2: PERCENTAGE OF STUDY POPULATION WITH
USG EVIDENCE OF FATTYLIVER
AND NO FATTY LIVER**




FIGURE 3: PERCENTAGE OF STUDY POPULATION HAVING EVIDENCE OF FATTY LIVER IN BOTH GROUPS.




FIGURE 4: PERCENTAGE OF POPULATION WITH NO EVIDENCE OF HEPATIC STEATOSIS IN ULTRASONOGRAPHY






FIGURE 5:- PERCENTAGE OF STUDY POPULATION WITH HIGH, INTERMEDIATE AND LOW RISK IN OBESE GROUP.



**FIGURE 6 : STAGE OF FIBROSIS IN FIBRO SCAN
IN OBESE GROUP.**



DISCUSSION

6.DISCUSSION

The study was conducted in government stanley medical college hospital and the study population was 100 patients. Informed consent was obtained from all the patients included in the study.

All the patients included were Type 2 Diabetes Mellitus patients diagnosed by WHO criteria. Study population was subjected to blood investigations, ultrasonography and selected patients for fibro scan. The data was collected and interpreted and analysed using IBM.SPSS statistics software 23.0 Version. To describe about the data descriptive statistics frequency analysis, percentage analysis were used for categorical variables.

The study population included 40 males and 60 females. Study population categorised to obese and non obese group. 57 % of population were in obese group and 43 % in non obese group. 59.6% of patients had ultrasonographic evidence of hepatic steatosis in obese group and 40.4% had no ultrasonographic evidence of hepatic steatosis in obese group.

6.9% of patients only had ultrasonographic evidence of hepatic steatosis in non obese group and 93% had no evidence of hepatic steatosis in non obese group. This shows the significance of body mass index and

obesity playing a major role in pathogenesis of non alcoholic fatty liver disease. The percentage of patients with fatty liver is significantly higher in obese group than in non obese group.

The non alcoholic fatty liver disease fibrosis score was calculated in all patients of study population and 35% of patients had high risk score in obese group compared to only 0.2% in non obese group.

36% of patients had intermediate risk in obese group compared to 13.9% in non obese group. 28% had low risk in obese group compared to 83.7% in non obese group. This showed that obese group had significantly higher risk of hepatic fibrosis compared to non obese group in diabetic patients.

When the patients of high risk NAFLD score of both groups including 20 patients from obese group and 1 patient from non obese group were subjected to fibro scan, the correlation between fibro scan and NAFLD score was similar and hence prove that non invasive tests have upcoming role in clinical practise compared to invasive tests like liver biopsy etc.. in early diagnosis and appropriate management for the group of patients with risk factors of non alcoholic fatty liver disease.

SUMMARY

7. SUMMARY

The study was done in 100 patients diagnosed to have Type 2 diabetes mellitus according to WHO criteria and categorised into obese and non obese groups according to Body Mass Index and ultrasonography was done to all subjects and categorised into patients with hepatic steatosis and no evidence of hepatic steatosis. Non alcoholic fatty liver disease fibrosis score was calculated for all patients and divided into three groups with low risk, intermediate and high risk.

Fibroscan was done for all patients with high risk NAFLD fibrosis score and correlation was studied, and all patients with high score was found to have significant stage of fibrosis according to METAVIR staging. Non alcoholic fatty liver disease fibrosis score was found significant in staging of hepatic fibrosis and correlated with fibro scan in staging of hepatic fibrosis or liver stiffness. NAFLD fibrosis score was found to be increased in obese patients, type 2 diabetes mellitus and helpful in early diagnosis of NAFLD and early intervention in reducing risk factors like hyperglycemia, hypertriglyceridemia, metabolic syndrome and reduce complications of NAFLD.

TABLES AND FIGURES

FIGURE 1: GENDER DISTRIBUTION IN STUDY POPULATION




FIGURE 2: AGE RANGES IN THE STUDY POPULATION.




TABLE 1 : MEAN , MEDIAN AND MODE FOR AGE:

OBSERVATIONS	TOTAL	MEAN	VARIANCE	STANDARD DEVATION
100.00	5083.00	50.830	108.80	10.43
MINIMUM	25%	MEDIAN	MAXIMUM	MODE
26.000	44.000	50.000	74.000	45%

**FIGURE 3: NAFLD FIBROSIS SCORE IN THE STUDY
POPULATION**




TABLE : 2 SEX DISTRIBUTION

SEX		
	Frequency	Percent
Male	40	40.0
Female	60	60.0
Total	100	100.0

**TABLE : 3: EVIDENCE OF FATTY LIVER IN OBESE
AND NON OBESE GROUP.**

	OBESE GROUP	NON OBESE GROUP
FATTY LIVER	34 (59.6%)	3 (6.9%)
NO EVIDENCE OF FATTY LIVER	23(40.4%)	40 (93.1%)
TOTAL	57	43

TABLE 4 : NAFLD SCORE IN OBESE AND NON OBESE GROUP

NAFLD SCORE	OBESE GROUP	NON OBESE GROUP
HIGH RISK	20 (35%)	1 (0.2%)
INTERMEDIATE RISK	21 (36 %)	6 (13.9%)
LOW RISK	16 (28%)	36 (83.7%)
TOTAL	57	43

TABLE : 5

	NAFLD SCORE			
AGE	1	2	3	Total
26	0	0	1	1
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	1.85%	1.00%
28	0	1	0	1
Row%	0.00%	100.00%	0.00%	100.00%
Col%	0.00%	4.00%	0.00%	1.00%
29	0	0	1	1
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	1.85%	1.00%
32	0	0	1	1
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	1.85%	1.00%
33	0	1	0	1
Row%	0.00%	100.00%	0.00%	100.00%
Col%	0.00%	4.00%	0.00%	1.00%
35	0	0	1	1
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	1.85%	1.00%

36	0	1	1	2
Row%	0.00%	50.00%	50.00%	100.00%
Col%	0.00%	4.00%	1.85%	2.00%
38	1	0	1	2
Row%	50.00%	0.00%	50.00%	100.00%
Col%	4.76%	0.00%	1.85%	2.00%
39	1	0	1	2
Row%	50.00%	0.00%	50.00%	100.00%
Col%	4.76%	0.00%	1.85%	2.00%
40	2	2	1	5
Row%	40.00%	40.00%	20.00%	100.00%
Col%	9.52%	8.00%	1.85%	5.00%
41	0	1	3	4
Row%	0.00%	25.00%	75.00%	100.00%
Col%	0.00%	4.00%	5.56%	4.00%
42	1	0	0	1
Row%	100.00%	0.00%	0.00%	100.00%
Col%	4.76%	0.00%	0.00%	1.00%
44	1	2	2	5
Row%	20.00%	40.00%	40.00%	100.00%
Col%	4.76%	8.00%	3.70%	5.00%

45	2	2	2	6
Row%	33.33%	33.33%	33.33%	100.00%
Col%	9.52%	8.00%	3.70%	6.00%
46	0	0	1	1
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	1.85%	1.00%
47	0	0	6	6
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	11.11%	6.00%
48	1	1	1	3
Row%	33.33%	33.33%	33.33%	100.00%
Col%	4.76%	4.00%	1.85%	3.00%
49	0	0	4	4
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	7.41%	4.00%
50	2	3	1	6
Row%	33.33%	50.00%	16.67%	100.00%
Col%	9.52%	12.00%	1.85%	6.00%
51	0	0	2	2
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	3.70%	2.00%

52	1	1	2	4
Row%	25.00%	25.00%	50.00%	100.00%
Col%	4.76%	4.00%	3.70%	4.00%
53	0	0	3	3
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	5.56%	3.00%
54	0	0	1	1
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	1.85%	1.00%
55	0	2	0	2
Row%	0.00%	100.00%	0.00%	100.00%
Col%	0.00%	8.00%	0.00%	2.00%
56	1	1	2	4
Row%	25.00%	25.00%	50.00%	100.00%
Col%	4.76%	4.00%	3.70%	4.00%
57	1	0	1	2
Row%	50.00%	0.00%	50.00%	100.00%
Col%	4.76%	0.00%	1.85%	2.00%
58	2	0	2	4
Row%	50.00%	0.00%	50.00%	100.00%
Col%	9.52%	0.00%	3.70%	4.00%

59	2	1	1	4
Row%	50.00%	25.00%	25.00%	100.00%
Col%	9.52%	4.00%	1.85%	4.00%
60	0	2	3	5
Row%	0.00%	40.00%	60.00%	100.00%
Col%	0.00%	8.00%	5.56%	5.00%
61	2	0	1	3
Row%	66.67%	0.00%	33.33%	100.00%
Col%	9.52%	0.00%	1.85%	3.00%
64	0	0	2	2
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	3.70%	2.00%
65	0	1	1	2
Row%	0.00%	50.00%	50.00%	100.00%
Col%	0.00%	4.00%	1.85%	2.00%
67	0	1	0	1
Row%	0.00%	100.00%	0.00%	100.00%
Col%	0.00%	4.00%	0.00%	1.00%
68	0	0	2	2
Row%	0.00%	0.00%	100.00%	100.00%

Col%	0.00%	0.00%	3.70%	2.00%
70	1	1	1	3
Row%	33.33%	33.33%	33.33%	100.00%
Col%	4.76%	4.00%	1.85%	3.00%
71	0	1	0	1
Row%	0.00%	100.00%	0.00%	100.00%
Col%	0.00%	4.00%	0.00%	1.00%
72	0	0	1	1
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	1.85%	1.00%
74	0	0	1	1
Row%	0.00%	0.00%	100.00%	100.00%
Col%	0.00%	0.00%	1.85%	1.00%
TOTAL	21	25	54	100
Row%	21.00%	25.00%	54.00%	100.00%
Col%	100.00%	100.00%	100.00%	100.00%

FREQUENCY TABLES

TABLE : 6

	Frequency	Percent
Obese	57	57.0
Non Obese	43	43.0
Total	100	100.0

TABLE : 7

USG

	Frequency	Percent
Fatty Liver	37	37.0
Non Fatty liver	63	63.0
Total	100	100.0

FREQUENCY TABLES

TABLE : 8

NAFLD SCORE

	Frequency	Percent
High risk	21	21.0
Intermediate risk	25	25.0
Low risk	54	54.0
Total	100	100.0

TABLE : 9

FIBRO SCAN

	Frequency	Valid Percent
F0 - F1	5	23.8
F2 - F3	13	61.9
F4	3	14.3
Total	21	100.0
Total	100	

TABLE 10

NAFLD SCORE * FIBRO SCAN Crosstabulation

	FIBRO SCAN			Total
	F0 - F1	F2 - F3	F4	
NAFLD SCORE High risk	5	13	3	21
Total	5	13	3	21

CONCLUSION

9.CONCLUSION

NAFLD is an increasingly important chronic liver disease next to alcoholic liver disease ranging from Steatosis,steatohepatitis to cirrhosis. Insulin resistance and oxidativestress play important roles in etiopathogenesis of non alcoholic liver disease. NAFLD is usually asymptomatic and incidentally diagnosed on routine laboratory investigation. Liver biopsy remains the most sensitive and specific means of providing prognostic information but nowadays other noninvasive tests are upcoming and playing a major role in clinical practise. In the absence of definite therapies, treatment is generally directed at optimizingbody weight and controlling risk factors. Liver transplantation is a therapeutic option for decompensated chronicliver disease.

ANNEXURE - I

MASTER CHART

MASTER CHART

S.NO	NAME	AGE	SEX	BMI	USG	NAFL D SCOR E	FIBRO SCAN
1	Amutha	58	2	1	1	1	3
2	Sivagami	50	2	2	1	2	
3	Muniammal	50	2	1	2	2	
4	Shanthi	48	2	1	2	2	
5	Razia	47	2	2	2	3	
6	Bhagyam	61	2	1	1	1	2
7	Krishnaveni	60	2	2	2	3	
8	Sherifa	49	2	2	2	3	
9	Maheshwari	50	2	1	1	1	2
10	Kousia Begum	44	2	1	2	3	
11	Sujatha	38	2	2	2	3	
12	Beevi john	70	2	1	1	2	
13	Kala	35	2	2	2	3	
14	Elizabeth	40	2	1	1	1	2
15	Mani	60	1	1	2	3	
16	arumugam	70	1	2	1	1	1
17	narayanan	64	1	2	2	3	
18	shanmugam	60	1	1	1	2	
19	thiyagarajan	59	1	1	1	1	2
20	vasudevan	49	1	2	2	3	
21	robert	71	1	1	1	2	
22	bhoopalan	44	1	2	2	2	
23	srinivasan	61	1	1	1	1	2
24	pachiappan	45	1	2	2	3	
25	anand	49	1	1	2	3	
26	rajendiran	47	1	2	2	3	

27	Kuppu	40	2	1	1	1	2
28	suseela devi	58	2	2	2	3	
29	arun	28	1	1	1	2	
30	chinnaponnu	65	2	1	2	2	
31	nagaraj	39	1	1	1	1	2
32	krishnan	56	1	1	1	1	3
33	mahalakshmi	44	2	2	2	3	
34	ahmed	52	1	1	2	3	
35	vasanthi	53	2	2	2	3	
36	raman	44	1	1	1	2	
37	meenai	36	2	2	2	2	
38	selvi	47	2	2	2	3	
39	asirvatham	48	1	1	2	3	
40	balaraman	65	1	2	2	3	
41	logeshwari	39	2	2	2	3	
42	sarawathy	56	2	1	2	3	
43	vasuki	48	2	1	1	1	2
44	sukuna	51	2	1	2	3	
45	nagaraj	41	1	2	2	3	
46	selvam	50	1	2	2	3	
47	udaiyappan	41	1	1	1	2	
48	padmavathy	70	2	2	2	3	
49	rajeshwari	58	2	1	2	3	
50	valli	54	2	2	2	3	
51	kamalam	55	2	1	1	2	
52	selvi	47	2	2	2	3	
53	meenambal	60	2	2	2	2	
54	amudha	40	2	1	1	2	
55	mariammal	45	2	1	2	3	
56	devaki	52	2	2	2	3	
57	jothi	42	2	1	1	1	2
58	chellammal	61	2	2	2	3	

59	kuppuraj	49	1	2	2	3	
60	rajan	45	1	1	1	2	
61	kannan	51	1	1	2	3	
62	krishnamoort hy	57	1	1	1	1	2
63	mariappan	64	1	2	2	3	
64	janaki	68	2	2	2	3	
65	perumal	56	1	1	2	3	
66	damodharan	72	1	2	2	3	
67	deivanai	74	2	2	2	3	
68	sarasu	58	2	1	1	1	2
69	kothaiaamal	68	2	2	2	3	
70	nancy	33	2	1	2	2	
71	kayalvizhi	44	2	1	1	1	1
72	kaniappan	56	1	1	1	2	
73	nivedha	29	2	2	2	3	
74	divya	38	2	1	1	1	1
75	mohan	53	1	1	2	3	
76	monisha	26	2	2	2	3	
77	murugan	45	1	1	1	1	1
78	govindammal	45	2	1	1	2	
79	rosammal	59	2	1	1	2	
80	pappu	50	2	2	2	2	
81	krishnamoort hy	41	1	2	2	3	
82	devi	50	2	1	1	1	2
83	kannan	36	1	1	2	3	
84	valliyammal	53	2	2	2	3	
85	rajan	59	1	1	1	1	3
86	sumathy	40	2	1	2	3	
87	murali	47	1	2	2	3	
88	senthil	52	1	1	1	1	2

89	kanchana	45	2	1	1	1	1
90	priya	32	2	2	2	3	
91	ramasamy	40	1	1	1	2	
92	neelakandan	47	1	1	2	3	
93	joselin	41	2	1	2	3	
94	monica	55	2	2	1	2	
95	ayesha begum	67	2	1	2	2	
96	thamaraiselvi	60	2	1	2	3	
97	ramachandra n	59	1	1	2	3	
98	ponni	46	2	2	2	3	
99	megala	52	2	1	1	2	
100	vimala	57	2	2	2	3	
			SEX	BMI	USG		NAFLD SCORE
			MALE- 1	OBES E-1	FATT Y LIVE R-1		HIGH RISK-1
			FEMAL E-2	NON- OBES E-2	NO FATT Y LIVE R-2		INTERMEDI ATE RISK-2
							LOW RISK-3

ANNEXURE - II

BIBLIOGRAPHY

BIBLIOGRAPHY

1. Falck-Ytter Y, Younossi Z M, Marchesini G, Mc Cullough AJ. Clinical features and natural history of nonalcoholic steatosis syndromes. *Semin Liver Dis* 2001;21:17-26.
2. Shimada M, Hashimoto E, Tanai M, Hasegawa K, Okuda H, Hayashi N, et al. Hepatocellular carcinoma in patients with non-alcoholic steatohepatitis. *J Hepatol* 2002; 37:154-60.
3. Ludwig J, Viggiano T, McGill D, Ott B. Nonalcoholic steatohepatitis. Mayo Clinic experience with a hitherto unnamed disease. *Mayo Clin Proc* 1980;55:434-8.
4. Neuschwander-Tetri BA, Caldwell SH. Nonalcoholic Steatohepatitis: Summary of an AASLD single topic conference. *Hepatology* 2003;37:1202-1219.
5. Sanyal AJ; American Gastroenterological Association. AGA technical review on nonalcoholic fatty liver disease. *Gastroenterology* 2002;123:1705-1725
6. Angulo P. Nonalcoholic fatty liver disease. *N Eng J Med* 2002;346:1221-1231.
7. Duseja A, Chawla Y. Nonalcoholic fatty liver disease in India: how much? How soon? *Trop Gastroenterol.* 2005;26:1-3. *Indian J Gastroenterol* (2010) 29:217-225 223

8. Singh SP, Nayak S, Swain M, et al. Prevalence of non-alcoholic fatty liver disease in coastal eastern India: a preliminary ultrasonographic survey. *Indian J Gastroenterol*. 2004;25:76–9.
9. Matteoni CA, Younossi ZM, Gramlich T, et al. Non alcoholic fatty liver disease: a spectrum of clinical and pathological severity. *Gastroenterology* 1999;116:1413–19.
10. Ekstedt M, Franzen LE, Mathiesen UL, et al. Long-term follow-up of patients with NAFLD and elevated liver enzymes. *Hepatology* 2006;44:865–73.
11. Wong VW, Wong GL, Choi PC, et al. Disease progression of non-alcoholic fatty liver disease: a prospective study with paired liver biopsies at 3 years. *Gut* 2010;59:969–74.
12. Fassio E, Alvarez E, Dominguez N, et al. Natural history of non alcoholic steatohepatitis: a longitudinal study of repeat liver biopsies. *Hepatology* 2004;40:820–6.
13. Adams LA, Sanderson S, Lindor KD, et al. The histological course of nonalcoholic fatty liver disease: a longitudinal study of 103 patients with sequential liver biopsies. *J Hepatol* 2005;42:132–8.
14. Matteoni CA, Younossi ZM, Gramlich T, et al. Non-alcoholic fatty liver disease: a spectrum of clinical and pathological severity. *Gastroenterology* 1999;116:1413–19.

15. Sanyal AJ, Banas C, Sargeant C, et al. Similarities and differences in outcomes of cirrhosis due to non-alcoholic steatohepatitis and hepatitis C. *Hepatology* 2006;43:682–9.
16. Ascha MS, Hanouneh IA, Lopez R, et al. The incidence and risk factors of hepatocellular carcinoma in patients with nonalcoholic steatohepatitis. *Hepatology* 2010;51:1972–8.
17. Anstee QM, Targher G, Day CP. Progression of NAFLD to diabetes mellitus, cardiovascular disease or cirrhosis. *Nat Rev Gastroenterol Hepatol* 2013;10:330–44.
18. Browning JD, Szczepaniak LS, Dobbins R, et al. Prevalence of hepatic steatosis in an urban population in the United States: impact of ethnicity. *Hepatology* 2004;40:1387–95.
19. Angulo P, Keach JC, Batts KP, et al. Independent predictors of liver fibrosis in patients with nonalcoholic steatohepatitis. *Hepatology* 1999;30:1356–62.
20. Musso G, Gambino R, De Michieli F, et al. Dietary habits and their relations to insulin resistance and postprandial lipemia in nonalcoholic steatohepatitis. *Hepatology* 2003;37:909–16.
21. Ouyang X, Cirillo P, Sautin Y, et al. Fructose consumption as a risk factor for non-alcoholic fatty liver disease. *J Hepatol* 2008;48:993–9.

22. Cortez-Pinto H, Jesus L, Barros H, et al. How different is the dietary pattern in non-alcoholic steatohepatitis patients? *Clin Nutr* 2006;25:816–23.
23. Molloy JW, Calcagno CJ, Williams CD, et al. Association of coffee and caffeine consumption with fatty liver disease, nonalcoholic steatohepatitis, and degree of hepatic fibrosis. *Hepatology* 2012;55:429–36.
24. Musso G, Cassader M, Olivetti C, et al. Association of obstructive sleep apnoea with the presence and severity of non-alcoholic fatty liver disease. A systematic review and metaanalysis. *Obes Rev* 2013;15:417–31.
25. Tatsumia K, Saibarab T. Effects of obstructive sleep apnoea syndrome on hepatic steatosis and non-alcoholic steatohepatitis. *Hepatol Res* 2005;33:100–4.
26. Sookoian S, Castano GO, Burgueno AL, et al. A nonsynonymous gene variant in the adiponutrin gene is associated with nonalcoholic fatty liver disease severity. *J Lipid Res* 2009;50:2111–6.
27. Romeo S, Kozlitina J, Xing C. Genetic variation in PNPLA3 confers susceptibility to nonalcoholic fatty liver disease. *Nat Genet* 2008;40:1461–5.

28. Angulo P. Nonalcoholic fatty liver disease. *N Eng J Med* 2002;346: 1221-1231.
29. Chitturi S, Farrell GC. Etiopathogenesis of nonalcoholic steatohepatitis. *Semin Liver Dis* 2001;21:27-41.
30. Day C, James O. Steatohepatitis: a tale of two “hits”? (editorial). *Gastroenterology* 1998;114:842-5.
31. Tilg H, Diehl AM. Cytokines in alcoholic and nonalcoholic steatohepatitis. *Eng J Med* 2000; 343:1467-1476.
32. Duseja A, Murlidharan R, Bhansali A, Sharma S, Das A, Das R, Chawla Y. Assessment of insulin resistance and effect of metformin in nonalcoholic steatohepatitis- A preliminary report. *Indian J Gastroenterol* 2004;23:12-15.
33. Blendis L, Oren R, Halpern Z. NASH: can we iron out the pathogenesis? *Gastroenterology* 2000; 118:981-5.
34. Duseja A, Das R, Nanda M, Das A, Grewal G, Chawla Y. Nonalcoholic steatohepatitis in Asian Indians is neither associated with iron overload nor with HFE gene mutations. *World J Gastroenterol* (in Press).
35. Uygun A, Kadayifci A, Yesilova Z, Erdil A, Yaman H, Saka M, Deveci MS, Bagci S, Gulsen M, Karaeren N, Dagalp K. Serum

- leptin levels in patients with nonalcoholic steatohepatitis. *Am J Gastroenterol* 2000 Dec;95:3584-9.
36. Wigg AJ, Thomson Roberts IC, Dymock RB, McCarthy PJ, Grose RH, Cummins AG. The role of small intestinal bacterial overgrowth, intestinal permeability, endotoxaemia, and tumor necrosis factor alfa in the pathogenesis of nonalcoholic steatohepatitis. *Gut* 2001;48:206-11.
37. Hotamisligil GS, Arner P, Caro JF, Atkinson RL, Spiegelman BM. Increased adipose tissue expression of tumor necrosis factor in human obesity and insulin resistance. *J Clin Invest* 1995;95:2409-2415.
38. Uysal KT, Wiesbrock SM, Marino MW, Hotamisligil GS. Protection from obesity -induced insulin resistance in mice lacking TNF- α function. *Nature* 1997;389:610-614.
39. Sanyal AJ, Sargent - Campbell C, Mirshahi F, Rizzo WB, Contos MJ, Sterling RK, et al. Nonalcoholic steatohepatitis: Association of insulin resistance and mitochondrial abnormalities. *Gastroenterology* 2001; 120: 1183-92.
40. Byrne CD. Ectopic fat, insulin resistance and non-alcoholic fatty liver disease. *Proc Nutr Soc* 2013;72:412–419.

41. Kantartzis K, Machann J, Schick F, Fritsche A, Haring HU, Stefan N. The impact of liver fat vs. visceral fat in determining categories of prediabetes. *Diabetologia* 2010;53:882–889.
42. Gao Z, Zhang J, Kheterpal I, Kennedy N, Davis RJ, Ye J. Sirtuin 1 (SIRT1) protein degradation in response to persistent c-Jun N-terminal kinase 1 (JNK1) activation contributes to hepatic steatosis in obesity. *J Biol Chem* 2011;24:22227–22234.
43. Samuel VT, Liu ZX, Qu X, Elder BD, Bilz S, Befroy D, et al. Mechanism of hepatic insulin resistance in non-alcoholic fatty liver disease. *J Biol Chem* 2004;30:32345–32353.
44. Byrne CD. Dorothy Hodgkin Lecture 2012: non-alcoholic fatty liver disease, insulin resistance and ectopic fat: a new problem in diabetes management. *Diabet Med* 2012;29:1098–1107.
45. Mehal WZ. The Gordian Knot of dysbiosis, obesity and NAFLD. *Nat Rev Gastroenterol Hepatol* 2013;10:637–644.
46. Jornayvaz FR, Shulman GI. Diacylglycerol activation of protein kinase C ϵ and hepatic insulin resistance. *Cell Metab* 2012;15:574–584.
47. Samuel VT, Shulman GI. Mechanisms for insulin resistance: common threads and missing links. *Cell* 2012;148:852–871.
48. Yamaguchi K, Yang L, McCall S, Huang J, Yu XX, Pandey SK, et al. Inhibiting triglyceride synthesis improves hepatic steatosis but

- exacerbates liver damage and fibrosis in obese mice with nonalcoholic steatohepatitis. *Hepatology* 2007;45:1366–1374.
- 49.Schenk S, Saberi M, Olefsky JM. Insulin sensitivity: modulation by nutrients and inflammation. *J Clin Invest* 2008;118:2992–3002.
50. Leroux A, Ferrere G, Godie V, Cailleux F, Renoud ML, Gaudin F, et al. Toxic lipids stored by Kupffer cells correlates with their pro-inflammatory phenotype at an early stage of steatohepatitis. *J Hepatol* 2012;57:141–149.
- 51.Perry RJ, Samuel VT, Petersen KF, Shulman GI. The role of hepatic lipids in hepatic insulin resistance and type 2 diabetes. *Nature* 2014;510:84–91. [66] Watt MJ, Barnett AC, Bruce CR, Schenk S, Horowitz JF, Hoy AJ. Regulation of plasma ceramide levels with fatty acid oversupply: evidence that the liver detects and secretes de novo synthesised ceramide. *Diabetologia* 2012;55:2741–2746.
- 52.Yang G, Badeanlou L, Bielawski J, Roberts AJ, Hannun YA, Samad F. Central role of ceramide biosynthesis in body weight regulation, energy metabolism, and the metabolic syndrome. *Am J Physiol Endocrinol Metab* 2009;297:E211–E224.
- 53..Kolesnick RN, Kronke M. Regulation of ceramide production and apoptosis. *Annu Rev Physiol* 1998;60:643–665.[69] Lipina C,

- Hundal H. Sphingolipids: agents provocateurs in the pathogenesis of insulin resistance. *Diabetologia* 2011;1:1596–1607.
54. Cantley JL, Yoshimura T, Camporez JP, Zhang D, Jornayvaz FR, Kumashiro N, et al. CGI-58 knockdown sequesters diacylglycerols in lipid droplets/ER preventing diacylglycerol-mediated hepatic insulin resistance. *Proc Natl Acad Sci U S A* 2013;110:1869–1874.
55. Hotamisligil GS. Endoplasmic reticulum stress and the inflammatory basis of metabolic disease. *Cell* 2010;140:900–917.
56. Tilg H, Moschen AR. Insulin resistance, inflammation, and non-alcoholic fatty liver disease. *Trends Endocrinol Metab* 2008;19:371–379.
57. Targher G, Byrne CD. Nonalcoholic fatty liver disease: a novel cardiometabolic risk factor for type 2 diabetes and its complications. *J Clin Endocrinol Metab* 2013;98:483–495.
58. Ix JH, Sharma K. Mechanisms linking obesity, chronic kidney disease, and fatty liver disease: the roles of fetuin-A, adiponectin, and AMPK. *J Am Soc Nephrol* 2010;21:406–412.
59. Marceau P, Biron S, Hould FS, Marceau S, Simard S, Thung SN, et al. Liver pathology and metabolic syndrome X in severe obesity. *J Clin Endocrinol Metab* 1999; 84:1513-17.

- 60.Dixon JB, Bhathal PS, O' Brien PE. Nonalcoholic fatty liver disease: Predictors of nonalcoholic steatohepatitis and liver fibrosis in the severely obese. *Gastroenterology* 2001;121:91-100.
- 61.Duseja A, Das A, Dhiman RK, Bhansali A, Chawla Y. Nonalcoholic steatohepatitis - Is it part of metabolic syndrome. *Indian J Gastroenterol* 2004;23:(Suppl 1):S16.
- 62.Perseghin G, Lattuada G, De CF, Esposito A, Belloni E, Ntali G, et al. Increased mediastinal fat and impaired left ventricular energy metabolism in young men with newly found fatty liver. *Hepatology* 2008;47:51–58.
- 63.Lautamaki R, Borra R, Iozzo P, Komu M, Lehtimaki T, Salmi M, et al. Liver steatosis coexists with myocardial insulin resistance and coronary dysfunction in patients with type 2 diabetes. *Am J Physiol Endocrinol Metab* 2006;291:E282–E290.
- 64.Rijzewijk LJ, Jonker JT, van der Meer RW, Lubberink M, de Jong HW, Romijn JA, et al. Effects of hepatic triglyceride content on myocardial metabolism in type 2 diabetes. *J Am Coll Cardiol* 2010;56:225–233.
- 65.Rijzewijk LJ, van der Meer RW, Smit JW, Diamant M, Bax JJ, Hammer S, et al. Myocardial steatosis is an independent predictor of diastolic dysfunction in type 2 diabetes mellitus. *J Am Coll Cardiol* 2008;52:1793–1799.

66. Sinner MF, Wang N, Fox CS, Fontes JD, Rienstra M, Magnani JW, et al. Relation of circulating liver transaminase concentrations to risk of newonset atrial fibrillation. *Am J Cardiol* 2013;111:219–224.
67. Alonso A, Misialek JR, Amiin MA, Hoogeveen RC, Chen LY, Agarwal SK, et al. Circulating levels of liver enzymes and incidence of atrial fibrillation: the Atherosclerosis Risk in Communities cohort. *Heart* 2014;100:1151–1156.
68. Algra A, Tijssen JG, Roelandt JR, Pool J, Lubsen J. QTc prolongation measured by standard 12-lead electrocardiography is an independent risk factor for sudden death due to cardiac arrest. *Circulation* 1991;83:1888–1894.
69. Robbins J, Nelson JC, Rautaharju PM, Gottdiener JS. The association between the length of the QT interval and mortality in the Cardiovascular Health Study. *Am J Med* 2003;115:689–694.
70. Yasui K, Sumida Y, Mori Y, Mitsuyoshi H, Minami M, Itoh Y, et al. Nonalcoholic steatohepatitis and increased risk of chronic kidney disease. *Metabolism* 2011;60:735–739.
71. Machado MV, Goncalves S, Carepa F, Coutinho J, Costa A, Cortez-Pinto H. Impaired renal function in morbid obese patients with nonalcoholic fatty liver disease. *Liver Int* 2012;32:241–248

72. Williams AL, Hoofnagle JH. Ratio of serum aspartate to alanine aminotransferase in chronic hepatitis. Relationship to cirrhosis. *Gastroenterology* 1988;95:734–9.
73. Shah AG, Lydecker A, Murray K, et al. Comparison of noninvasive markers of fibrosis in patients with non-alcoholic fatty liver disease. *Clin Gastroenterol Hepatol* 2009;7:1104–12.
74. Sterling RK, Lissen E, Clumeck N, et al. Development of a simple noninvasive index to predict significant fibrosis in patients with HIV/HCV coinfection. *Hepatology* 2006;43:1317–25.
75. Adams LA, George J, Bugianesi E, et al. Complex non-invasive fibrosis models are more accurate than simple models in non-alcoholic fatty liver disease. *J Gastroenterol Hepatol* 2011;26:1536–43.
76. McPherson S, Anstee QM, Henderson E, et al. Are simple noninvasive scoring systems for fibrosis reliable in patients with NAFLD and normal ALT levels? *Eur J Gastroenterol Hepatol* 2013;25:652–8.
77. Sandrin L, Fourquet B, Hasquenoph JM, et al. Transient elastography: a new noninvasive method for assessment of hepatic fibrosis. *Ultrasound Med Biol* 2003;29:1705–13.

78. Foucher J, Chanteloup E, Vergniol J, et al. Diagnosis of cirrhosis by transient elastography (FibroScan): a prospective study. *Gut* 2006;55:403–8.
79. Wong VW, Vergniol J, Wong GL, et al. Diagnosis of fibrosis and cirrhosis using liver stiffness measurement in non-alcoholic fatty liver disease. *Hepatology* 2010;51:454–62.
80. Castera L, Foucher J, Bernard PH, et al. Pitfalls of liver stiffness measurement: a 5-year prospective study of 13,369 examinations. *Hepatology* 2010;51:828–35.
81. Dahl JJ, Pinton GF, Palmeri ML, et al. A parallel tracking method for acoustic radiation force impulse imaging. *IEEE Trans Ultrason Ferroelectr Freq Control* 2007;54:301–12.

ANNEXURE - III

PROFORMA

PROFORMA

1. NAME:

2. AGE:

3. SEX:

4. BMI:

5. DIABETES MELLITUS:

6. PLATELET COUNT:

7. ASPARTATE TRANSAMINASE:

8. ALANINE TRANSAMINASE:

9. SERUM ALBUMIN:

10.FORMULA:

$-1.675 + 0.037 * \text{AGE (years)} + 0.094 * \text{BMI (kg/m}^2\text{)} + 1.13 * \text{IFG/}$
 $\text{DIABETES (yes =1,no=0)} + 0.99 * \text{AST/ALT ratio} - 0.013 * \text{platelet (*}$
 $10^9 /\text{l)} - 0.66 * \text{albumin (g/dl)}.$

10. ULTRASOUND:

FATTY LIVER- YES/NO

CONSENT FORM

STUDY TITLE : “NONALCOHOLIC FATTY LIVER DISEASE FIBROSIS SCORE IN DIABETIC PATIENTS”

STUDY CENTRE : Govt. Stanley Medical College, Chennai-1

PARTICIPANT NAME : AGE: SEX: IP/OP NO:

I confirm that I have understood the purpose of procedure for the above study, I have the opportunity to ask the question and all my questions and doubts have been answered to my satisfaction.

I understand that the investigator, regulatory authorities and the ethics committee will not need my permission to look at my health records both in respect to the current study and any further research that may be conducted in relation to it, even if I withdraw from the study. I understand that my identity will not be revealed in any information released to third parties or published, unless as required under the law. I agree not to restrict the use of any results that arise from the study.

I hereby consent to participate in this study of : **“NONALCOHOLIC FATTY LIVER DISEASE FIBROSIS SCORE IN DIABETIC PATIENTS”**

.

Signature of Investigator:

Place :

Date :

Study Investigators Name

Institution

Signature / Thumb Impression of patient